

**SARS-COV-2/ INFLUENZA A+B/RSV / ADENOVIRUS/
M.PNEUMONIAE ANTIGEN COMBO RAPID TEST
(NASOPHARYNGEALSWAB /NASOPHARYNGEAL
WASH/NASAL ASPIRATE)- COMBO KIT (6IN1)**



**Get a Fast and Easy Solution for Confusable Diseases
Screening with CITEST Rapid Tests**

Comparison of Common Symptoms

Symptom	COVID-19	Flu	RSV	Adeno	MP	Cold
Fever	√	√	√	√	√	√
Cough	√	√	√	√	√	√
Headache	√	√			√	
Running nose		√	√			√
Stuffy nose		√				√
Sneeze			√			√
Sore throat	√	√	√	√	√	√
Breath shortness	√		√		√	
Sputum	√			√	√	
Vomit		√		√		
Diarrhea		√		√		
Weak limbs	√					
Muscle ache		√				

* Due to disease progression and individual differences, some symptoms may not appear.


Test Sensitivity, Specificity and Accuracy

Analytes	Sensitivity	Specificity	Accuracy
SARS-CoV-2	96.4%	99.0%	98.2%
Flu A	95.0%	99.1%	98.4%
Flu B	92.9%	99.1%	98.1%
RSV	94.3%	96.2%	95.9%
Adeno	96.9%	98.6%	98.4%
MP	91.8%	98.0%	97.0%

* Test has been evaluated with specimens obtained from the patients. RT-PCR is used as the reference method.


COUNTRY OF ORIGIN : CANADA

CAT NO.
IRT-CT545
SPECIMEN
NASOPHARYNGEALSWAB/NASOPHARYNGEAL WASH/NASAL ASPIRATE
PACK
20TEST/PER BOX
READING TIME
15 MINUTES



SPECIMEN COLLECTION, TRANSPORT AND STORAGE

1. Insert a sterile swab into the nostril of the patient, reaching the surface of the posterior nasopharynx.
2. Swab over the surface of the posterior nasopharynx 5-10 times.
3. Withdraw the sterile swab from the nasal cavity and avoid excess volume and highly-viscous nasopharyngeal discharge.



Specimen Transport and Storage

Specimens should be tested as soon as possible after collection. If swabs are not been processed immediately, it is highly recommended the swab specimen is placed into a dry, sterile, and tightly sealed plastic tube for storage. The swab specimen in dry and sterile condition is stable for up to 24 hours at 2-8 °C.

+60 12-684 2078

sales@vigagemilang.com.my

B-G-8, RAWANG IDAMAN 1,

JALAN IDAMAN 1,

TAMAN RAWANG IDAMAN, 48000 RAWANG SELANGOR