

C. difficile Rapid Test

***C. difficile* Rapid Test** is a qualitative immunoassay for detection of *Clostridium difficile* Glutamate Dehydrogenase (GDH), Toxin A and Toxin B in human fecal specimens.



About *C. difficile* infection (CDI)

Clostridium difficile (*C. difficile*) is the major causative agent of diarrhea and colitis associated with antibiotics.

CDI is considered responsible for approximately 25% of the diarrhea incidents related to the consumption of antibiotics, and CDI can lead to pseudomembranous colitis (PMC), requiring urgent treatment with antibiotics effective against *C. difficile* and which, without treatment, may severely compromise the life of patients. In the case of PMC, CDI mortality can be as high as 6% to 30%. Rapid immunoassay of CDI has become more important.

Notable Features

- High-quality & trusted diagnostic products
- Instant screening test
- Simplified sampling & procedure
- Rapid, sensitive, and accurate results



B-G-8 RAWANG IDAMAN, JALAN IDAMAN 1, TAMAN RAWANG
IDAMAN,
48000 RAWANG SELANGOR DARUL EHSAN

Phone Number : 012 - 313 3081 Email: sales@vigagemilang.com.my

Product Information

Information Type	Specimen	Detection	Format	Package	Time to results	Shelf life & Storage Conditions	Qualification
<i>C. difficile</i> Toxin A/B Rapid Test Device	Feces	<i>Clostridium difficile</i> Toxin A, Toxin B in human feces	Multi-Window Device	25 pcs/box	10 minutes	24 months at 2-30 °C	CE
<i>C. difficile</i> Toxin A/B & GDH Rapid Test Device	Feces	<i>Clostridium difficile</i> Toxin A, Toxin B and glutamate dehydrogenase (GDH) antigen in human feces	Multi-Window Device	20 pcs/box	10 minutes	24 months at 2-30 °C	CE
<i>C. difficile</i> Toxin A Rapid Test Device	Feces	<i>Clostridium difficile</i> Toxin A in human feces	Device	25 pcs/box	10 minutes	24 months at 2-30 °C	CE
<i>C. difficile</i> Toxin B Rapid Test Device	Feces	<i>Clostridium difficile</i> Toxin B in human feces	Device	25 pcs/box	10 minutes	24 months at 2-30 °C	CE
<i>C. difficile</i> GDH Rapid Test Device	Feces	<i>Clostridium difficile</i> glutamate dehydrogenase (GDH) antigen in human feces	Device	25 pcs/box	10 minutes	24 months at 2-30 °C	CE

Test Procedure

